

A task and performance analysis of endoscopic submucosal dissection (ESD) surgery

Berk Cetinsaya¹ · Mark A. Gromski² · Sangrock Lee³ · Zhaohui Xia³ · Doga Demirel¹ · Tansel Halic⁴ · Coskun Bayrak⁵ · Cullen Jackson⁶ · Suvarnu De³ · Sudeep Hegde⁶ · Jonah Cohen⁶ · Mandeep Sawhney⁶ · Stavros N. Stavropoulos⁷ · Daniel B. Jones⁶

Received: 15 February 2018 / Accepted: 10 August 2018
© Springer Science+Business Media, LLC, part of Springer Nature 2018

Abstract

Background ESD is an endoscopic technique for en bloc resection of gastrointestinal lesions. ESD is a widely-used in Japan and throughout Asia, but not as prevalent in Europe or the US. The procedure is technically challenging and has higher adverse events (bleeding, perforation) compared to endoscopic mucosal resection. Inadequate training platforms and lack of established training curricula have restricted its wide acceptance in the US. Thus, we aim to develop a Virtual Endoluminal Surgery Simulator (VESS) for objective ESD training and assessment. In this work, we performed task and performance analysis of ESD surgeries.

Methods We performed a detailed colorectal ESD task analysis and identified the critical ESD steps for lesion identification, marking, injection, circumferential cutting, dissection, intraprocedural complication management, and post-procedure examination. We constructed a hierarchical task tree that elaborates the order of tasks in these steps. Furthermore, we developed quantitative ESD performance metrics. We measured task times and scores of 16 ESD surgeries performed by four different endoscopic surgeons.

Results The average time of the marking, injection, and circumferential cutting phases are 203.4 (σ : 205.46), 83.5 (σ : 49.92), 908.4 s. (σ : 584.53), respectively. Cutting the submucosal layer takes most of the time of overall ESD procedure time with an average of 1394.7 s (σ : 908.43). We also performed correlation analysis (Pearson's test) among the performance scores of the tasks. There is a moderate positive correlation ($R=0.528$, $p=0.0355$) between marking scores and total scores, a strong positive correlation ($R=0.7879$, $p=0.0003$) between circumferential cutting and submucosal dissection and total scores. Similarly, we noted a strong positive correlation ($R=0.7095$, $p=0.0021$) between circumferential cutting and submucosal dissection and marking scores.

Conclusions We elaborated ESD tasks and developed quantitative performance metrics used in analysis of actual surgery performance. These ESD metrics will be used in future validation studies of our VESS simulator.

✉ Tansel Halic
tanselh@uca.edu

¹ Department of Computer Science, University of Arkansas at Little Rock, Little Rock, AR, United States

² Division of Gastroenterology and Hepatology, Indiana University School of Medicine, Indianapolis, IN, USA

³ Rensselaer Polytechnic Institute, Troy, NY, United States


⁴ Department of Computer Science, University of Central Arkansas, 201 Donaghey Ave, Conway, AR 72035, USA

⁵ Department of Computer Science & Information Systems, Youngstown State University, Youngstown, OH, USA

⁶ Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA, USA

⁷ NYU Winthrop Hospital, New York, NY, USA

Graphical abstract

**Keywords** Endoscopic training · Endoscopic submucosal dissection · Colorectal cancer · ESD

The risk of developing colorectal cancer during the lifetime is 1 in 21 for men and 1 in 23 for women [1]. According to the World Center Research Fund International [2], with an average of 1.4 million new cases per year, colorectal cancer is the third most commonly encountered cancer type in the world. In the US, it is the second leading cause of death, with 14.8 deaths per 100,000 men and women according to the reports published by National Cancer Institute [3]. The estimated new colon cancer diagnoses in 2017 is 135,430, which is equivalent to 8% of all new cancer diagnoses. The estimated colon cancer-related death in 2017 is 50,260 people [3]. Unless timely steps in terms of screening, diagnosis, and treatment are taken, colorectal cancer will continue to be a major health challenge to the US and the world.

There are several techniques for the management of premalignant colorectal polyps and superficial colorectal cancer (i.e., not penetrating the submucosa). For mucosal-based lesions, the most widely used technique in the US currently is endoscopic mucosal resection (EMR). En bloc resection can often be accomplished for lesions less than 20 mm [4] with EMR. For lesions larger than 20 mm or for non-standard shaped lesions (i.e., over a fold), complete removal often requires piecemeal resection. Piecemeal resection can obscure resection borders on the pathologic specimen and can make a pathological diagnosis of an R0 resection difficult. Therefore, ESD was developed in order to dissect larger lesions (> 20 mm) with an en bloc resection, allowing for more complete pathological review [5–9].

In general, ESD has a longer procedure time compared to EMR [4, 10]. Compared with EMR, ESD has higher rates

of procedural complications (bleeding and perforation) [11]. While it causes higher risks of perforation, ESD has a lower rate of delayed bleeding [6], higher en bloc resection rate, and lower local recurrence rate [10, 12]. Furthermore, due to the en bloc resection of the lesion, it is easier for a complete pathological analysis of the tumor by the pathologist [13].

ESD for colorectal tumors is a complex procedure due to the relatively thin colonic wall and difficulties keeping the position of the endoscope stable due to the shape, folds, and contractions of the colon [14]. Given the relative difficulty of ESD compared to EMR, an expert endoscopist is required to have extensive training in the procedure. This study is motivated by the established need for a standardized method of training and evaluating colorectal ESD.

ESD training in the United States

Training in ESD has been difficult to coordinate for practitioners in the US. In Asia, most endoscopists start ESD by observing and performing ESD of superficial gastric tumors. Given the dramatic differences in incidence of gastric cancer and resultant differences in gastric cancer screening policies, endoscopists in the US generally do not encounter many superficial gastric cancers in clinical practice [15]. In addition, guidelines from US-based GI and endoscopy societies provide little guidance [16].

Conventional training methods for ESD are mostly based on animal models [17–20] and patient-based observed and proctored training. Fujishiro et al. [21] derived a

labor-intensive proposal of training for ESD with multiples steps. Ex-vivo simulators pose some problems with translatability (difficulty with electrosurgical settings in desiccated specimens, inability to simulate intraprocedural bleeding) and set-up. While in-vivo simulations with live animal models [22] are realistic, they are not widely used due to ethical concerns and associated cost. With the use of animal models, there is a learning curve of 15–30 cases for gastric ESD [18, 22]. Given the difficulties in training noted above, there is tremendous opportunity for another modality of ESD training in the United States.

Alternative training methods such as virtual reality (VR)-based simulators may be an effective adjunct to a training regimen for ESD. A high-fidelity VR simulator would allow for skills training in both the advanced psychomotor and cognitive skills aspects of ESD. The simulator can be used to train not only endoscopists but also other operating room (OR) personnel (endoscopy nurses and technicians) in all of procedural aspects [23]. The advantages of endoscopy-based VR simulators can be summarized as, (1) no ethical dilemma with initial training through an apprentice model on human subjects, with no risk of harm to the patient or malpractice, since no-human subject is involved [24]; (2) repeatable on different tumor-types and sizes; (3) affordable compared to recurrent costs of other animate and inanimate training models [25, 26]; (4) potentially reduced training time with a steeper learning curve once human ESD work commences [27]; (5) and quantifiable measurement for both assessment and training [28].

VR simulators have become widespread in the last decade to improve psychomotor learning outcomes in laparoscopic surgery and natural orifice transluminal endoscopic surgery (NOTES) [28, 29]. However, there is no virtual reality-based simulation for colorectal ESD. The VR-development for ESD is challenging, given the constricted and detailed endoscopic operating space, large deformation of tissue, complicated modeling (e.g., various tumor models), and complex physics-based interactions (e.g., incision, cutting, dissection, etc.) between different kinds of endoscopic accessories and colon tissue [28].

Our ultimate goal is to design and develop a high-fidelity VR-based colorectal ESD simulator using haptics technology. We aim to create a teaching and assessment platform for structured ESD training with objective feedback. We hypothesize that this platform could significantly improve ESD training and equip trainees with a high level of proficiency in the ESD steps, which would prepare them for initial proctored training in humans. In order to achieve this, the aim of this current study is to perform a hierarchical task analysis (HTA) of the ESD procedure, derive metrics for quantitative performance measurement of actual endoscopic steps, and to perform a time analysis of the actual ESD procedures in an attempt to develop authentic measures.

Materials and methods

Task analysis methods

HTA, in the context of an overall procedural analysis, is a method that details the steps of a procedural process from beginning to end. The specific aim of this hierarchical expression is to identify crucial details in each step of the procedure and allow the creation of metrics for each step. These details will then be translated and programmed into the VR simulation platform. We performed a thorough literature review of the colorectal ESD procedure and had discussions with expert endoscopists from the US, Japan, and Korea during the analysis. As a result, we detailed the ESD phases and tasks performed in each phase. The task tree, the systematic representation of the tasks and their order, and objective grading metrics were created as part of our task analysis.

Grading metrics

Based on the tasks that were discovered in the HTA, we developed grading metrics for each task and subtask of each ESD phase with the input of expert endoscopists. We utilized a Likert-like scale for task scoring. The highest score point of 3–5 is given for the best/optimal action. A suboptimal action is given a point value of 1–3, depending on the importance and consequence of the action. A trainee receiving a 0 score in a task signifies that no proper action is taken. In some metrics (such as in bleeding management criteria), a 0 score denotes an overall failing of the training exercise. This is termed a “kill switch” for the training procedure. The grading system was derived by consensus from expert endoscopists.

Time and performance analysis

Based on the HTA, we created timing guidelines with specified actions for “start” and “end” for each specified task and subtask. This was created by expert consensus. Sixteen videos of ESD procedures performed by four different endoscopists were analyzed. In the time analysis of the ESD videos, three raters reviewed each video individually and independently using VLC and Windows media players. All raters were also debriefed with the evaluation guidelines including possible rare cases. It was noted that multiple subtasks (such as hemostasis or injection) could be recurrent events in multiple phases of the procedure. The raters were trained to time these tasks within duration of phases. Several segments or clips of the ESD videos were timed and scored with raters prior to the study to increase the rater training.

Once raters confirmed that they have a clear understanding about HTA and all ESD steps in detail and timing and scoring guidelines, they timed all phases' and subtasks' start and end time and rated scores (based upon scoring criteria devised in Results section “[Grading metrics](#)”) for all the videos. We performed inter-rater reliability (IRR) for all phases using Fleiss' kappa [30, 31] to evaluate the agreement among raters.

Prior to our study, we hypothesized that there may be relation between the task duration and scores from the devised grading metrics. This stems from the common observation of relation noted in the literature that skills and experience level of the surgeon affects the total surgery time, time spent in a task and performance score [32, 33]. In order to quantify the relation, we performed Pearson's correlation tests between times and scores. We used in *RStudio* version 1.0.153 with R version 3.4.1 for the statistical analysis.

Results

Task analysis

Six major phases for the ESD procedure were identified: (1) procedural preparation, (2) coagulative marking circumferentially outside of the borders of the lesion, in order to improve visualization of the boundaries of the tumor during the procedure, (3) injection of a solution into the submucosal space to lift the lesion and create a protective cushion [34] for cutting, (4) circumferential cutting around the lesion using endoscopic electrocautery knives which have been described elsewhere (Matsui et al. [35]), (5) submucosal dissection by using an electrosurgical knife, and (6) evaluation of the colon for bleeding and perforations throughout the procedure and at the conclusion of the procedure. Furthermore, we described all the tasks performed for each of six ESD phases. We determined the necessary tasks and optional/selective tasks for each phase. The hierarchy of the phases and execution order of tasks form the hierachal task tree. The tasks trees from preparation to evaluation phase are found in Figs. 1, 2, 3, 4, 5 and 6. In addition to the phases, we created task trees for perforation (Fig. 7) and bleeding management (Fig. 8) which are commonly performed sets of tasks but not phases per se. In these tasks trees, normal processes are illustrated in rectangle and decisions are shown in diamonds. Arrows show the progression of the tasks in the order presented and they should be performed in order until the completion of the tasks in the tree. Some surgical tasks are recurrent where the tasks should be repeated until the task's goal is met. These repetitive tasks are illustrated with an arrow that directs the execution order back to the former steps.

Preparation phase

The preparation step (Fig. 1) is the first step of ESD. Before the procedure, tools and injection solutions must be prepared. The type of the endoscope is selected according to the location of the lesion; an upper esophagogastrroduodenoscope may be used if the lesion is located in the left colon or rectum, whereas a pediatric or adult colonoscope is most likely used if it is located in the right colon [12]. In the US, all ESD procedures are performed at a minimum with procedural moderate sedation and often with general anesthesia depending on the availability of the anesthesiologist, practice-, and procedure-specific considerations (e.g., estimated duration of procedure, estimated difficulty of procedure, co-morbidities of the patient). In ESD, a carbon dioxide (CO₂) insufflation system is used to reduce the patient's pain from colonic gas retention and risk of peritoneal distention should a perforation occur during the procedure. Furthermore, a clear endoscopic cap is used at the distal end of the scope. This helps to facilitate traction while dissecting the submucosal space. Finally, the patient will be positioned ideally so that the lesion will be in an anti-gravity position for increasing the lifting and elevation during the dissection and to aid in dissection [5, 36–39].




Fig. 1 Steps in preparation phase

Marking phase

The marking phase in the task tree is illustrated in Fig. 2. In the beginning of the marking step, the selected scope is inserted through the anus and navigated through the colon until the lesion is located. There are four potential methods for detecting and confirming the aspects and pathological components of the lesion: (1) high-definition white light, (2) narrow band imaging, (3) near focus/magnification, and (4) chromoendoscopy. The marking step could be completely skipped if the lesion is clear and visible after chromoendoscopy. Depending on the case, a physician may choose to utilize multiple methods. After successful detection and evaluation, the next step is to clean the lesion and surrounding mucosa with a water jet. In some cases, small debris may be present and must be removed with the suction function of the endoscope. Thereafter, the knife, depending on the

endoscopist's preference, is inserted into the working channel of the scope. Then, the electrosurgical unit (ESU) should be confirmed to be set for the desired tissue effect and at the discretion of the expert. The recommended settings for each ESD phase specific to ESU can be found in [5]. After setting the ESU, the cutting surface/blade of the knife is exposed in the colonoscopy view.

Although there are various techniques for marking the mucosal layer, two techniques are very common; the first technique for marking is performed by placing the knife onto the surface of the mucosa and creating coagulation marks approximately 2–3 mm apart and 5 mm peripheral to the lesion. The second technique is to use an argon plasma coagulation (APC) probe in close proximity to mucosa and use the PULSED APC mode on the ESU.

Injection phase

Based on the primary ESD knife being used, the injection phase will vary (Fig. 3). If the endoscopist is not using a multi-purpose knife such as the HybridKnife, then the current marking tool (knife or APC probe) is retracted and removed from the working channel of the scope and an injection needle is inserted into the working channel. After the injection needle catheter is present into the colonic lumen, the needle tip is exposed, and the needle tip is inserted into the submucosal layer. Injection is performed until sufficient elevation is achieved. After sufficient elevation is achieved, the needle tip is retracted. Note that it is not necessary to perform last two steps if the instrument is a dual-purpose knife. By using a multi-purpose knife, injection of the lifting




Fig. 2 Steps in marking phase




Fig. 3 Steps in injection phase

solution can be performed at any time without switching instruments.

Circumferential cutting phase

The first step in this phase is inserting the selected knife into the working channel of the scope (see Fig. 4). Then, the ESU should be set to the correct specification. Once the cutting mode and power settings in the ESU are adjusted, the cutting surface/blade of the knife is exposed. The next step is to perform the marginal incision around the lesion until the submucosal fibers are exposed. This is accomplished by first penetrating through the mucosa (in a lesion already injected with submucosal injection) while cutting; this is performed carefully so as not to damage the underlying muscle layer—this is the initial cut. After this first incision is made through the mucosa, a cut around the circumference of the lesion is made. Oftentimes, the circumferential cut will be performed in stages (i.e., not entirely circumferential initially) so as to preserve the submucosal cushion for a longer period of time. Generally, this cutting is performed just outside the perimeter of the previous markings, with care not to cut into the lesion itself. Throughout the circumferential cut, the depth of cutting should be monitored, to ensure cutting is not too deep (into muscle) or too shallow (not entirely through mucosa). A high quality circumferential cut will help the efficiency of the later steps of the ESD. In the case

of inadequate lesion elevation, the injection step should be repeated. When bleeding occurs, the site is examined, washed and, if necessary, hemostasis is performed.

Submucosal dissection phase

Submucosal dissection (Fig. 5) is the most critical step of the ESD procedure. Inserting the knife, setting the ESU and exposing the knife's tip are performed the same as in the previous steps. Then, the submucosal layer is dissected by manipulating the knife below the lesion. Again, similar to the previous phase, the injection step should be repeated until elevation is sufficient and also cleaning or hemostasis is repeated when bleeding occurs. Otherwise, dissection is advanced until the lesion separates from the underlying colon and is completely resected. The approach to the lesion will need to be altered throughout the dissection phase. It is important that all cutting is performed within the submucosal plane, preferably closer to the muscle side than the mucosa side, with avoidance of cutting the muscle. Small and precise movements are of paramount




Fig. 4 Steps in circumferential cutting phase




Fig. 5 Steps in submucosal dissection phase

importance. During the dissection, small veins and arteries will be encountered penetrating the submucosal space. These should be prophylactically managed with either the coagulation setting of the knife or with the coagrasper device. The clear endoscopic cap is used to facilitate traction of the submucosal space while cutting. After complete resection, the lesion is removed with appropriate graspers, or by suction into the endoscopic cap. Although it is not common, a snare can be used as a salvage technique to remove the lesion if dissection becomes tedious.

Evaluation phase

The last phase of the ESD is a colonoscopic review (see Fig. 6) performed in order to detect possible perforations, residual lesion, or visible vessels after the removal of lesion. When perforation (see Fig. 7) occurs, CO_2 should be decreased and consideration for peritoneal decompression should ensue. Often with large perforations, the intraluminal colon will no longer insufflate even with regular flow. Bleeding (see Fig. 8) is treated with endo-clips or hemostasis tools (e.g., coagrasper, APC, bipolar probe). At the end of the procedure, pinning the specimen on a mounting board allows for numerous objectives to be met in addition to analyzing the dimensions of the resection, including allowing for proper pathological fixation without distortion and also to examine the surface of the lesion for en bloc status.




Fig. 6 Steps in evaluation phase




Fig. 7 Steps in perforation management

Grading metrics

Metrics for each task and subtask in the HTA and grading schema were determined by expert consensus (Table 1). There are some common and repeated metrics in miscellaneous scores such as bleeding intervention times, address of knife angle to the dissection plane, knife handling, cleaning blood from the field, etc. These tasks need to be performed during most of the phases rather than a specific phase or task. For some of these metrics, the criteria might occur multiple times such as clearing the visibility or hemostasis times. In these cases, the endoscopist will be assessed only once and the minimum score will be recorded. For example, if the endoscopist requires more than 60 s for hemostasis while using coagraspers or doing coagulation hemostasis, or more than 120 s for hemostasis by hemoclip in one hemostasis scenario, s/he will receive one point. In addition, there are no acceptable metrics




Fig. 8 Steps in bleeding management

in timing performance in interventional flexible endoscopy available in the literature. The subject matter experts determined timing metrics based on expert opinion.

Time and performance analysis

The specified timing guidelines were created for all phases and subtasks (see Table 2) by expert consensus. These specify criteria when raters need to start and end time for each phase and tasks. This is needed to avoid ambiguity and eliminate inconsistency among the videos and reviewers.

Sixteen ESD videos were reviewed. These videos are internal colonoscopy recordings of live ESD procedures except one that was performed on a porcine colon. In only five videos, the “Removing the lesion” subtask was recorded. Similarly, the “Evaluation” phase was only recorded in five videos. In one of the videos, the marking phase was not recorded. In these cases, the timings and scores for these tasks were not considered in the statistical analysis. All videos started at the marking phase; therefore, time analysis and scores are not available for the preparation phase.

According to time analysis performed, the submucosal dissection phase is the longest phase with maximum time variation (see Table 3; Fig. 9). The average time of the marking phase (see Fig. 10) includes detection of the lesion, marking the lesion, washing the lesion, and spraying the dye times (if performed). Circumferential cutting has three subtasks (see Fig. 11): cleaning the bleeding, hemostasis, injection tasks. We determined that the resolution of the bleeding task time takes longer time in the circumferential cutting phase compared to the submucosal dissection time (See Figs. 9, 10).

We also computed scores with respect to our ESD metrics presented in Table 3. These scores were computed by

Table 1 Grading metrics for all main phases and subtasks

Preparation	
Position of patient	
Lesion is in an anti-gravity position	3
Lesion is not in an anti-gravity position	0
Selection of the endoscope	
Correct endoscope	3
Incorrect endoscope	0
Status of the distal cap attachment	
Clear distal cap	3
No distal cap	0
Marking	
Insertion of HD endoscope	
Navigate to the lesion	2
Advance beyond the lesion	1
Stopping proximal to the lesion	0
Identification of the lesion within 60 s	
Identified	3
Not identified	0
Cleaning the lesion and surrounding mucosa by water jet	
Washed satisfactorily	3
There is still debris	1
If there is small debris around the colon	
Suctioned completely	3
Not suctioned completely	1
Marking—distance between each mark	
2–3 mm (ideal margin)	5
3–5 mm (acceptable)	3
0–1 mm or > 5 mm (poor margin)	1
No marking	0
Peripheral zone of lesion marking	
5–7 mm	5
More than 7 mm	3
Less than 5 mm	1
Knife position into mucosa on marking step	
Close to mucosa (in PULSED APC mode) or 0.5 mm inside to mucosa (in soft coagulation mode)	5
Less than 0.5 mm (soft coagulation mode)	3
Greater than 0.5 mm (soft coagulation mode)	1
Chromoendoscopy is used	
Spraying enough dye	5
Spraying insufficient dye	1
Injection	
Injection of solution	
Enough mL solution	5
Too much solution	3
Not enough solution	1
Not injected any solution	0 (fail)
Lesion elevation	
Sufficient elevation (enough fluid cushion)	5
Not enough elevation (not enough fluid cushion)	3
Circumferential cutting and submucosal dissection	

Table 1 (continued)

Choosing knife type	
Correct knife	5
Incorrect tool ^a	0 (fail)
Initial incision	
Start circumferential cutting 1–2 mm outer distal proximity of marking points	5
Start circumferential cutting greater than 2 mm outer distal proximity of marking points	3
Start circumferential cutting on marking points	1
Start circumferential cutting 1–2 mm inner distal proximity of marking points	0 (fail)
Dissection	
Cutting middle zone or muscle zone of the submucosal layer	5
Cutting while not having adequate cushion	3
Cutting the muscle (muscularis propria) layer	1
Cutting the mucosa layer inadvertently	0
Perforation of the lesion during the dissection	
No perforation	5
Perforation	0
Resecting the lesion	
En bloc resection	5
Piecemeal complete resection	3
Incomplete resection	0
Removing the lesion	
Lesion successfully removed	3
Failure to remove the lesion	0
Evaluation	
Colonoscopic review	
Detecting perforations	5
Not detecting perforations	0
Miscellaneous	
Bleeding intervention time ^b	
Less than 60 s (coagrasper, doing coagulation hemostasis) or 120 s (hemoclip)	3
More than 60 s (coagrasper, doing coagulation hemostasis) or 120 s (hemoclip)	1
No intervention	0 (fail)
Perforation: avoidance	
Injecting more solution	3
Not injecting more solution	0
Visible bleeding location	
Hemostasis	5
No hemostasis	0 (fail)
Non-visible bleeding location	
Injecting solution	5
Not injection	0
Cleaning the blood	
Spray the water	3
Not spray the water	0
Knife handling	
Smoothness and gentleness in tool handling	5

Table 1 (continued)

Discrete motions in tool handling	3
Aggressive tool handling	0
Knife angle to the dissection plane	
15–35° with respect to the dissection surface (except IT knife)	3
Degree [5–15] or [35–45] with respect to the dissection surface	1
Other angles	0 (fail)
Knife exposing	
Expose the knife on accurate step	3
Expose the knife on another step	0
Knife retracting	
Retract the knife on accurate step	3
Retract the knife on another step	0
Position of HD endoscope	
Near the lesion	2
Far from the lesion	1
Tasks execution order	
Completion of tasks executed in order	2
Completion of tasks executed not in order	0

^aThis metric is attributed to instrument choices made that is not relevant for a specific phase

^bThe subject matter experts determined timing metrics based on expert opinion

the same independent raters who performed the task timing of the videos. Based on our results, the average score of the videos is 79.3 (max: 93.8 points, min: 49.0 points) (see Fig. 12). The majority of the attainable scores belong to the category of the circumferential cutting and submucosal dissection scores and miscellaneous scores. The miscellaneous scores are mostly referred to the tasks that can be performed at any phase (e.g., cleaning the blood, tool handling, identifying bleeding location) so they do not fall under one specific phase of the surgery.

We computed the IRR for 3 raters of the 16 videos (4 phases evaluated in videos) for agreement in scores. The agreement rate of the preparation phase is $\kappa=1.00$. There is a substantial agreement ($\kappa=0.69$) for the marking phase. The discrepancy in scores are primarily due to the difficulty to assess the exact marking distance (in mm) from the videos. There is excellent agreement ($\kappa=1.00$) for the injection phase. For the circumferential cutting and submucosal dissection phases, there is a substantial agreement ($\kappa=0.76$). For the miscellaneous scores, the agreement rate is an excellent agreement ($\kappa=0.84$). In general, we conclude that the raters have a high level of inter-rater reliability on the average scores for the ESD tasks.

We categorized Pearson's correlation test results in time-score correlation (see Table 4) and score-score correlation of phases (see Table 5). Our results demonstrate that the

Table 2 Start and end times

Phase	Tasks
Marking	Insertion of the HD endoscope Insertion of the HD endoscope into anus Seeing the lesion Detection of the lesion Seeing the lesion Exposing the knife Washing the lesion and surrounding mucosa Starting flushing water Stopping flushing water Suction the debris Starting suction of debris Stopping suction of the debris Spraying dye Insertion of the spraying tool Finishing spraying dye Marking the margin of the lesion Insertion of the knife into mucosa Completion of the circumferential marking around the lesion Cleaning the bleeding if occurs Start spraying water Stop spraying water Hemostasis if bleeding occurs Insertion of the hemostasis tool into the submucosa Removing the tool from the submucosa Injection
Injection	Insert the injection needle into the endoscope Retracting the injection needle
Circumferential cutting	Cutting the mucosal layer Insertion of the knife into the mucosal layer surrounding the lesion Removing the knife from the mucosal layer when circumferential incision is completed Cleaning the bleeding if occurs Spraying water Stop flushing Hemostasis if bleeding occurs Insertion of the hemostasis tool into the submucosa Removing the tool from the submucosa Injection Same as previous injection step Same as previous injection step

Table 2 (continued)

Phase	Tasks
Submucosal dissection	Cutting the submucosal layer
	Insertion of the knife into the submucosal layer surrounding the lesion
	Resection of the lesion
	Cleaning the bleeding if occurs
	Spraying water
	Stop flushing
	Hemostasis if bleeding occurs
	Insertion of the hemostasis tool into the submucosa
	Removing the tool from the submucosa
	Injection
	Same as previous injection step
	Same as previous injection step
	Removing the lesion
	Insertion of the grasping tool
	Removing from the endoscope
	Colonoscopic review
Evaluation	Identification of the colon wall around the resected area
	Removing the endoscope from colon
	Cleaning the bleeding if occurs
	Spraying water
	Stop flushing
	Hemostasis if bleeding occurs
	Insertion of the hemostasis tool into the submucosa
	Removing the tool from the submucosa

Table 3 Time analysis of phases

Phases	Avg. (s)	Min (s)	Max (s)	σ (s)
Marking	203.4	23	885	205.46
Injection	83.5	24	212	49.92
Circumferential cutting	908.4	301	2390	584.53
Subtasks				
Cleaning the bleeding	24.9	1	97	29.76
Hemostasis	28.8	5	75	22.98
Injection	104.7	8	245	75.04
Submucosal dissection	1394.7	75	3196	908.43
Subtasks				
Cleaning the bleeding	15.1	1	41	14.03
Hemostasis	28.4	1	90	26.70
Injection	117.6	15	454	128.73
Total time of phases	2327.6	856	6297	1399.28
Total time of videos	3049.8	1020	9553	2021.20

majority of the task times and task scores are negatively related.

Discussion

In this study, we performed a hierarchical task analysis and task trees to determine the individual integral steps to the colonic ESD procedure. Furthermore, we produced a scoring metric to evaluate colonic ESD procedures. Finally, we independently evaluated colonic ESD procedure videos to determine the timing of each component of the HTA and to administer the scoring metric for these procedures.

In our scoring metrics, higher points in Likert scale are given for the optimal or desired actions that are designated to reflect better surgery performance (see “[Grading metrics](#)”). Negative correlations in our results (Tables 4, 5) convey that the endoscopists who have higher scores in our metrics generally complete the procedure in a shorter period of time than the ones with lower scores. This may be attributed to (a) the skill level of surgeon or (b) the difficulty of the specific lesion, that can result in more rapid task completion times overall.

We identified that there is a strong positive correlation between miscellaneous scores and total scores, circumferential cutting and submucosal dissection scores and total scores. Endoscopists who have a high score from

Fig. 9 Time analysis of submucosal dissection phase




Fig. 10 Time analysis of marking phase





Fig. 11 Time analysis of circumferential cutting phase



miscellaneous tasks tend to have a high score from circumferential cutting and submucosal dissection tasks as well.

One limitation of the study is that we analyzed a total of 16 videos from four endoscopists. It is possible that validating the scoring metric with a more extensive study that involves more endoscopists and more procedures would

provide more robust procedural data. Furthermore, colorectal ESD is a dynamic field, with frequent modifications to the tools available and techniques employed. There can be many variations including various assistive devices for traction or colonic stability, or dissection strategies similar to the pocket technique.

Fig. 12 Average scores of all phases




Table 4 Time—score correlations

Time	Score	R	p	Correlation
Cleaning the bleeding in circumferential cutting	Marking phase	-0.6159	0.0252	Negative moderate
Cleaning the bleeding in submucosal dissection	Marking phase	-0.6824	0.0298	Negative moderate
Detection of the lesion on marking phase	Circumferential cutting and submucosal dissection	-0.6921	0.0061	Moderate negative
Cutting the mucosal layer in circumferential cutting phase	Circumferential cutting and submucosal dissection	-0.6323	0.0086	Moderate negative
Detection of the lesion in marking phase	Total	-0.6439	0.0131	Moderate negative
Marking phase	Total	-0.5139	0.0505	Negative moderate
Cutting the submucosal layer in submucosal dissection phase	Marking phase	0.0095	0	No correlation

Table 5 Score—score correlations

Score	Score	R	p	Correlation
Marking	Total	0.528	0.0355	Moderate positive
Circumferential cutting and submucosal dissection	Total	0.7879	0.0003	Strong positive
Miscellaneous	Total	0.8738	<0.00001	Strong positive
Circumferential cutting and submucosal dissection	Marking	0.7095	0.0021	Strong positive

In conclusion, we performed a HTA and developed a rubric for performance metrics for ESD. Based on the HTA and metrics, we carried out time and performance analysis of actual ESD videos. We presented correlations between task times and scores. We will integrate the steps and sub-steps of the ESD procedure and relative timing of each step (discovered in the procedural video analysis) as the foundation for the software and hardware development for our VR ESD simulator. Furthermore, the performance

metrics will be integrated into the simulation environment to provide the trainee with real-time feedback via quantitative performance measurement. The identified correlations will be compared with our simulator as a future work.

Acknowledgements This project was made possible by the Arkansas INBRE program, supported by a grant from the National Institute of General Medical Sciences, (NIGMS), P20 GM103429 from the National Institutes of Health. This project was also supported by National Institutes of Health (NIH) Grant NIH/NHLBI 1R01HL119248-01A1, NIH/NIBIB 2R01EB005807, 5R01EB010037, 1R01EB009362, R01CA197491 and 1R01EB014305.

Compliance with ethical standards

Disclosures Tansel Halic reports grants from National Institutes of Health, during the conduct of the study. Cullen Jackson reports grants from NIH via subcontract from RPI, during the conduct of the study. Suvrana De grants from National Institutes of Health, during the conduct of the study. Sudeep Hegde reports grants from National Institutes of Health, during the conduct of the study; other from State University of New York at Buffalo, outside the submitted work. Daniel B. Jones reports personal fees from Allurion, outside the submitted work. Berk Cetinsaya, Mark A. Gromski, Sangrock Lee, Zhaohui Xia, Doga Demirel, Coskun Bayrak, Jonah Cohen, Mandeep Sawhney, Stavros N. Stavropoulos have no conflicts of interest or financial ties to disclose.

References

- Key Statistics for Colorectal Cancer. <https://www.cancer.org/cancer/colon-rectal-cancer/about/key-statistics.html>. Accessed 14 Aug 2017
- Worldwide data | World Cancer Research Fund International. <https://www.wcrf.org/int/cancer-facts-figures/worldwide-data>. Accessed 28 May 2018
- Cancer of the Colon and Rectum—Cancer Stat Facts. <https://seer.cancer.gov/statfacts/html/colorect.html>. Accessed 13 Apr 2017
- Kantsevoy SV, Adler DG, Conway JD, Diehl DL, Farraye FA, Kwon R, Mamula P, Rodriguez S, Shah RJ, Song LMWK, Tierney WM (2008) Endoscopic mucosal resection and endoscopic submucosal dissection. *Gastrointest Endosc* 68:11–18. <https://doi.org/10.1016/j.gie.2008.01.037>
- Yamamoto K, Michida T, Nishida T, Hayashi T, Naito M, Ito T (2015) Colorectal endoscopic submucosal dissection: recent technical advances for safe and successful procedures. *World J Gastrointest Endosc* 7:1114–1128. <https://doi.org/10.4253/wjge.v7.i14.1114>
- Fujihara S, Mori H, Kobara H, Nishiyama N, Matsunaga T, Ayaki M, Yachida T, Morishita A, Izuishi K, Masaki T (2014) Current innovations in endoscopic therapy for the management of colorectal cancer: from endoscopic submucosal dissection to endoscopic full-thickness resection. *BioMed Res Int*. <https://doi.org/10.1155/2014/925058>
- Saito Y, Yamada M, So E, Abe S, Sakamoto T, Nakajima T, Otake Y, Ono A, Matsuda T (2014) Colorectal endoscopic submucosal dissection: technical advantages compared to endoscopic mucosal resection and minimally invasive surgery. *Dig Endosc Off J Jpn Gastroenterol Endosc Soc* 26(Suppl 1):52–61. <https://doi.org/10.1111/den.12196>
- Shono T, Ishikawa K, Ochiai Y, Nakao M, Togawa O, Nishimura M, Arai S, Nonaka K, Sasaki Y, Kita H (2011) Feasibility of endoscopic submucosal dissection: a new technique for En Bloc resection of a large superficial tumor in the colon and rectum. *Int J Surg Oncol* 2011:e948293. <https://doi.org/10.1155/2011/948293>
- Wang J, Zhang X-H, Ge J, Yang C-M, Liu J-Y, Zhao S-L (2014) Endoscopic submucosal dissection vs endoscopic mucosal resection for colorectal tumors: a meta-analysis. *World J Gastroenterol WJG* 20:8282–8287. <https://doi.org/10.3748/wjg.v20.i25.8282>
- Hotta K, Oyama T, Shinohara T, Miyata Y, Takahashi A, Kitamura Y, Tomori A (2010) Learning curve for endoscopic submucosal dissection of large colorectal tumors. *Dig Endosc* 22:302–306. <https://doi.org/10.1111/j.1443-1661.2010.01005.x>
- Oka S, Tanaka S, Kaneko I, Mouri R, Hirata M, Kawamura T, Yoshihara M, Chayama K (2006) Advantage of endoscopic submucosal dissection compared with EMR for early gastric cancer. *Gastrointest Endosc* 64:877–883. <https://doi.org/10.1016/j.gie.2006.03.932>
- Niimi K, Fujishiro M, Goto O, Kodashima S, Koike K (2012) Safety and efficacy of colorectal endoscopic submucosal dissection by the trainee endoscopists. *Dig Endosc* 24:154–158. <https://doi.org/10.1111/j.1443-1661.2012.01251.x>
- Sakamoto T, Takamaru H, Mori G, Yamada M, Kinjo Y, So E, Abe S, Otake Y, Nakajima T, Matsuda T, Saito Y (2014) Endoscopic submucosal dissection for colorectal neoplasms. *Ann Transl Med*. <https://doi.org/10.21037/3536>
- Kato H, Sakamoto T, Otsuka H, Yamada R, Watanabe K (2012) Colorectal cancer—from prevention to patient care. InTech
- Fukami N (2014) Endoscopic submucosal dissection, an issue of gastrointestinal endoscopy clinics, E-Book. Elsevier Health Sciences
- ASGE Standards of Practice Committee, Faulx AL, Lightdale JR, Acosta RD, Agrawal D, Bruining DH, Chandrasekhara V, Eloubeidi MA, Gurudu SR, Kelsey L, Khashab MA, Kothari S, Muthusamy VR, Qumseya BJ, Shaukat A, Wang A, Wani SB, Yang J, DeWitt JM (2017) Guidelines for privileging, credentialing, and proctoring to perform GI endoscopy. *Gastrointest Endosc* 85:273–281. <https://doi.org/10.1016/j.gie.2016.10.036>
- Gromski MA, Cohen J, Saito K, Gonzalez J-M, Sawhney M, Kang C, Moore A, Matthes K (2017) Learning colorectal endoscopic submucosal dissection: a prospective learning curve study using a novel ex vivo simulator. *Surg Endosc*. <https://doi.org/10.1007/s00464-017-5484-2>
- Kato M, Gromski M, Jung Y, Chuttani R, Matthes K (2013) The learning curve for endoscopic submucosal dissection in an established experimental setting. *Surg Endosc* 27:154–161. <https://doi.org/10.1007/s00464-012-2402-5>
- Parra-Blanco A, Arnau MR, Nicolás-Pérez D, Gimeno-García AZ, González N, Díaz-Acosta JA, Jiménez A, Quintero E (2010) Endoscopic submucosal dissection training with pig models in a Western country. *World J Gastroenterol WJG* 16:2895–2900. <https://doi.org/10.3748/wjg.v16.i23.2895>
- Berr F, Ponchon T, Neureiter D, Kiesslich T, Haringsma J, Kaehtler GF, Schmoll F, Messmann H, Yahagi N, Oyama T (2011) Experimental endoscopic submucosal dissection training in a porcine model: learning experience of skilled Western endoscopists. *Dig Endosc Off J Jpn Gastroenterol Endosc Soc* 23:281–289. <https://doi.org/10.1111/j.1443-1661.2011.01129.x>
- Fujishiro M (2009) Endoscopic submucosal dissection for colorectal neoplasms. *World J Gastrointest Endosc* 1:32. <https://doi.org/10.4253/wjge.v1.i1.32>
- Tsuji Y, Fujishiro M, Kodashima S, Niimi K, Ono S, Yamamichi N, Koike K (2014) Desirable training of endoscopic submucosal dissection: further spread worldwide. *Ann Transl Med*. <https://doi.org/10.3978/j.issn.2305-5839.2014.02.05>
- Sankaranarayanan G, Li B, Miller A, Wakily H, Jones SB, Schwartzberg S, Jones DB, De S, Olasky J (2016) Face validation of the virtual electrosurgery skill trainer (VEST®). *Surg Endosc* 30:730–738. <https://doi.org/10.1007/s00464-015-4267-x>
- Demirel D, Yu A, Halic T, Sankaranarayanan G, Ryason A, Spindler D, Butler KL, Cao C, Petrusa E, Molina M, Jones D, De S, Demoya M, Jones S (2016) Virtual airway skills trainer (VAST) simulator. *Stud Health Technol Inform* 220:91–97
- Qiao W, Bai Y, Lv R, Zhang W, Chen Y, Lei S, Zhi F (2014) The effect of virtual endoscopy simulator training on novices: a systematic review. *PLoS ONE* 9:e89224. <https://doi.org/10.1371/journal.pone.0089224>
- Triantafyllou K, Lazaridis LD, Dimitriadis GD (2014) Virtual reality simulators for gastrointestinal endoscopy training. *World J Gastrointest Endosc* 6:6–12. <https://doi.org/10.4253/wjge.v6.i1.6>
- Aggarwal R, Black SA, Hance JR, Darzi A, Cheshire NJW (2006) Virtual reality simulation training can improve inexperienced surgeons' endovascular skills. *Eur J Vasc Endovasc Surg Off J Eur Soc Vasc Surg* 31:588–593. <https://doi.org/10.1016/j.ejvs.2005.11.009>
- Pan JJ, Chang J, Yang X, Liang H, Zhang JJ, Qureshi T, Howell R, Hickish T (2015) Virtual reality training and assessment in laparoscopic rectum surgery. *Int J Med Robot Comput Assist Surg MRCAS* 11:194–209. <https://doi.org/10.1002/rcs.1582>
- Beyer-Berjot L, Berdah S, Hashimoto DA, Darzi A, Aggarwal R (2016) A virtual reality training curriculum for laparoscopic colorectal surgery. *J Surg Educ* 73:932–941. <https://doi.org/10.1016/j.jsurg.2016.05.012>
- Rücker G, Schimek-Jasch T, Nestle U (2012) Measuring inter-observer agreement in contour delineation of medical imaging in a dummy run using Fleiss' Kappa. *Methods Inf Med* 51:489–494. <https://doi.org/10.3414/ME12-01-0005>

31. Shrout PE, Fleiss JL (1979) Intraclass correlations: uses in assessing rater reliability. *Psychol Bull* 86:420–428. <https://doi.org/10.1037/0033-2909.86.2.420>
32. Demirel D, Butler KL, Halic T, Sankaranarayanan G, Spindler D, Cao C, Petrusa E, Molina M, Jones DB, De S, deMoya MA (2016) A hierarchical task analysis of cricothyroidotomy procedure for a virtual airway skills trainer simulator. *Am J Surg* 212:475–484. <https://doi.org/10.1016/j.amjsurg.2015.08.029>
33. Demirel D, Yu A, Cooper-Baer S, Dendukuri A, Halic T, Kockara S, Kockara N, Ahmadi S (2016) A hierarchical task analysis of shoulder arthroscopy for a virtual arthroscopic tear diagnosis and evaluation platform (VATDEP). *Int J Med Robot*. <https://doi.org/10.1002/rcs.1799>
34. Hsu W-H, Sun M-S, Lo H-W, Tsai C-Y, Tsai Y-J (2013) Clinical practice of endoscopic submucosal dissection for early colorectal neoplasms by a colonoscopist with limited gastric experience. *Gastroenterol Res Pract* 2013:e262171. <https://doi.org/10.1155/2013/262171>
35. Matsui N, Akahoshi K, Nakamura K, Ihara E, Kita H (2012) Endoscopic submucosal dissection for removal of superficial gastrointestinal neoplasms: a technical review. *World J Gastrointest Endosc* 4:123–136
36. Maple JT, Dayyeh BKA, Chauhan SS, Hwang JH, Komanduri S, Manfredi M, Konda V, Murad FM, Siddiqui UD, Banerjee S (2015) Endoscopic submucosal dissection. *Gastrointest Endosc* 81:1311–1325. <https://doi.org/10.1016/j.gie.2014.12.010>
37. Ribeiro MS, Wallace MB (2015) Endoscopic treatment of early cancer of the colon. *Gastroenterol Hepatol* 11:445–452
38. Shida T (2011) Endoscopic submucosal dissection for colorectal lesions. In: *Colonoscopy*, InTech. <https://doi.org/10.5772/20825>
39. Yamamoto H (2010) Endoscopic submucosal dissection for colorectal tumors. *Techn Gastroint Endosc* 27:287–295. <https://doi.org/10.1159/000258370>